Journal of Immunological Methods, 155 (1992) 273–274

© 1992 Elsevier Science Publishers B.V. All rights reserved 0022-1759/92/$05.00

Contents to volume 155 (1992)

Number 1 – 19 October 1992

Research reports

In vitro measurement and screening of monoclonal antibody affinity using fluorescence photobleaching
E.N. Kaufman and R.K. Jain (USA) 1

Optimization of cytotoxic assay by target cell retention of the fluorescent dye carboxyfluorescein diacetate (CFDA) and comparison with conventional ^{51}CR release assay
M. Provinciali, G. Di Stefano and N. Fabris (Italy) 19

Development of a new technique for recovery of cytokines from inflammatory sites in situ
R. Alam, T.C. Sim, K. Hilsmeier and J.A. Grant (USA) 25

Enzyme immunoassays for the estimation of adenosine 3′,5′ cyclic monophosphate and guanosine 3′,5′ cyclic monophosphate in biological fluids
J.K. Horton, R.C. Martin, S. Kalinka, A. Cushing, J.P. Kitcher, M.J. O'Sullivan and P.M. Baxendale (UK) 31

Stimulation of human T cells by sparse antigens captured on immunomagnetic particles
S. Hawke, N. Willcox, G. Harcourt, A. Vincent and J. Newsom-Davis (UK) 41

A peroxidase-independent method for the quantitation of extracellular hydrogen peroxide generated by activated phagocytes in vitro
R. Anderson (South Africa) 49

Immunological properties of ribosome-inactivating proteins and a saporin immunotoxin
P. Strocchi, L. Barbieri and F. Stirpe (Italy) 57

Use of anti-peptide antibodies for the design of antigen-specific immune complex assays
T. Maruyama, G.B. Thornton, S. Iino, K. Kurokawa and D.R. Milich (USA, Japan) 65

Molecular cloning of murine monoclonal anti-idiotypic Fab
Y. Kasai, D. Herlyn, M. Sperlagh, H. Maruyama, S. Matsushita and A.J. Linnenbach (USA, Japan) 77

Acoustic probe-based ELISA
D.K. Boraker, S.J. Bugbee and B.A. Reed (USA) 91

T helper epitopes enhance the cytotoxic response of mice immunized with MHC class I-restricted malaria peptides
C. Widmann, P. Romero, J.L. Maryanski, G. Corradin and D. Valmori (Switzerland) 95

A rapid isolation procedure for dendritic cells from mouse spleen by centrifugal elutriation
M.A. Ossevoort, R.E.M. Toes, M.L.H. De Bruijn, C.J.M. Melief, C.G. Figdor and W.M. Kast (Netherlands) 101

Rapid attachment of a helper T cell epitope to branched peptides by fragment condensation to give enhanced immunogenicity
G.W. McLean, A.M. Cross, M.S. Munns and H.S. Marsden (UK) 113

Restoring antibody activity of a monoclonal anti-RNP antibody by dissociative HPLC. Demonstration of blocking antibody binding sites with antigen released from effete hybridoma cells
J. Ma, B. Walsh, S.-L. Chen, R. Penny and S.N. Breit (Australia, China) 121

An improved method for the purification of IgG monoclonal antibodies from culture supernatants
D.A. Brooks, T.M. Bradford and J.J. Hopwood (Australia) 129

Effector-target interactions: saturability, affinity and binding isotherms. A study of such interactions in the human NK cell-K562 tumour cell system
P. Garcia-Peñarrubia, L. Cabrera, R. Alverez and J. Galvez (Spain) 133

Notes to authors 149

Number 1 – 5 November 1992

Research reports

Superoxide production by normal and chronic granulomatous disease (CGD) patient-derived EBV-transformed B cell lines measured by chemiluminescence-based assays
C.D. Porter, M.H. Parkar, M.K.L. Collins, R.J. Levinsky and C. Kinnon (UK) 151

The M20 IL-1 inhibitor. I. Purification by preparative isoelectric focusing in free solution
D. Peritt, I. Flechner, E. Okunev, P. Yanai, T. Halperin, A.J. Treves and V. Barak (Israel) 159
The M20 IL-1 inhibitor. II. Biological characterization
D. Peritt, I. Flechner, P. Yanai, E. Okunev, T. Halperin, A.J. Treves and V. Barak (Israel) 167
Antibodies to soluble human T cell receptor β chain recognize multiple epitopes on cell surface TcR
G.S. Basi, M.B. Riggs, K. Nash and R. Singer (USA) 175
Preparative-scale purification and characterization of MHC class II monomers
D. Passmore, D. Kopa and B. Nag (USA) 193
A complement-dependent enzyme immunoassay (C-EIA) with increased sensitivity for IgM-rich rabbit sera
H.-M. Kuhn, L. Brade and H. Brade (Germany) 201
Capture ELISA for IgM antibodies against *Plasmodium falciparum* glutamate rich protein
M. Dziegiel, M.B. Borre, E. Petersen, B. Högh, S. Jepsen, J. Vuust and N. Axelsen (Denmark) 207
A peroxidase-linked enzyme immunoassay for tumour necrosis factor α utilising alternative colorimetric or chemilumimetric substrates
W.R. Lamb, R.S.H. Pumphrey and P.E.C. Brenchley (UK) 215
An enzyme-linked immunosorbent assay for the quantification of solubilized CD14 in biological fluids
U. Grunwald, C. Krüger, J. Westermann, A. Lukowsky, M. Ehlers and C. Schütt (Germany) 225
Removal of false-positive reactions from plasma in an enzyme immunoassay for bovine interferon-γ
S.L. Jones, J.C. Cox, J.M. Shepherd, J.S. Rothel, P.R. Wood and A.J. Radford (Australia) 233
Analysis of glycoforms present in two mouse IgG2a monoclonal antibody preparations
J.M. Coco-Martin, F. Brunink, T.A.M. Van der Velden-de Groot and E.C. Beuvery (Netherlands) 241
An antibody capture bioassay (ACB) for DNase in human serum samples
U.E.M. Gibson, A.B. Chen, D.L. Baker and D.V. Sinicropi (USA) 249
Flow cytometric analysis of transport activity in lymphocytes electroporated with a fluorescent organic anion dye
J.E. Dinchuk, K.A. Kelley and G.N. Callahan (USA) 257
Use of the lipid emulsion system and *Salmonella typhimurium* mitogen adjuvant to stimulate IgG production in chickens
C. McCune, R.H. Goddard, S.-O. Cho and S.M. Wick (USA) 267

Book review 271

Announcement 272

Contents to volume 155 (1992) 273

Notes to authors 275